

# Pelvic Organ Prolapse

## What is a pelvic organ prolapse?

A pelvic organ prolapse occurs when there is a defect or weakness in the supporting structures of the pelvic organs. The pelvic floor muscle is a big hammock of muscles that supports the uterus, bladder and bowel. 1 in 9 women will develop a pelvic organ prolapse by the age of 80. There is a reported 29% recurrence rate of pelvic organ prolapse in women who have previous surgery for pelvic organ prolapse, with the surgical techniques of today this have been reduced.

## What causes it?

There are a number of contributing causes of prolapses:

- Childbirth or pregnancy esp. large babies >9lbs (4 kgs), forceps delivery
- Ageing and menopause
- Being overweight
- Chronic cough & chronic constipation
- Constant heavy lifting
- Previous pelvic surgery
- Collagen deficiency (collagen is a natural protein that helps keep tissues plump and elastic)

## How do I know I have one?


Common complaints with a prolapse are:

- Sensation of "something coming down" when up and about, disappears when lying down.
- Backache
- Increased frequency of needing to void {which is probably due to incomplete emptying of the bladder.}
- Problem emptying bowel
- A feeling of fullness or pressure in the lower abdominal area, {which is probably due to venous congestion and pressure from the abdominal contents on a weak and inadequate pelvic floor}.
- Sexual dysfunction and discomfort during intercourse.
- Urinary incontinence.


While pelvic organ prolapses are not life threatening, they can have a significant effect on overall wellbeing and quality of life.

## Types of Vaginal Prolapse

**CYSTOCELE (ANTERIOR WALL PROLAPSE):** A cystocele is when there is a defect in the anterior wall supports and causes, the bladder to prolapse into the vaginal wall. This contributes to inability to empty the bladder properly and stress urinary incontinence.



**RECTOCELE (POSTERIOR WALL PROLAPSE):** A rectocele is where the defect is in the posterior or back wall supports of the vagina and as a result the rectum bulges into the vagina. This can contribute to difficulty emptying the bowel as this causes a pocket to form and as a result faeces become lodged there, it is not uncommon for women with a rectocele to have to use perineal pressure to aid the emptying of their bowel. A rectocele can also cause a decreased stream when passing urine as the bulge presses up against the urethra (water pipe) thus obstructing the flow of urine.




# Pelvic Organ Prolapse



urogynaecology patient  
information

**ENTEROCOELE:** This is where the small bowel prolapses into the top of the vagina between the vagina and rectum, most commonly seen with a rectocele and or uterine prolapse.




**UTERINE PROLAPSE:** This is where the defect is in the structure that supports the womb, resulting in the cervix and uterus to prolapse into the vagina. There are varying degrees of this prolapse:

Grade 1: The uterus has dropped slightly and many women are unaware of the prolapse and are noticed during a routine pelvic check up.

Grade 2: The cervix and uterus has dropped further and can be felt or seen just inside the vagina.

Grade 3-4: The cervix and uterus can be seen or felt outside the vagina. This is the most severe of uterine prolapses and also known as a **procidentia**.




**VAGINAL VAULT PROLAPSE:** The vaginal vault is the top of the vagina.

With a vault prolapse this is where the vaginal wall loses its support from surrounding structure and the vagina falls in on itself. A vault prolapse can only occur to women who have had a previous hysterectomy. Vault prolapse statistics show increased risk of vault prolapse in women who had a hysterectomy for a uterine prolapse, the risk of this occurring are decreasing with new methods of surgery for prolapse repair. There are varying degrees of prolapses and are graded as:

Grade 1: prolapse is evident on examination but patient usually unaware of prolapse

Grade 2: Prolapse can be felt or seen at vaginal opening

Grade 3: Prolapse protrudes through the vaginal opening.



*The treatment of pelvic organ prolapse is assessed on an individual basis. Treatment may include pelvic muscle exercises, use of intravaginal pessaries, or surgical repair of the prolapse. Your specialist will discuss the best treatment options with you.*